

Potentially Preventable Hospitalization and Emergency Room Indicators

Hospitalization (Inpatient) Indicators
Hospitalization Rate due to Adult Asthma
Hospitalization Rate due to Asthma
Hospitalization Rate due to Asthma - Pediatric (ages 0-17)
Hospitalization Rate due to Community-Acquired Pneumonia
Hospitalization Rate due to COPD
Hospitalization Rate due to Dehydration
Hospitalization Rate due to Diabetes
Hospitalization Rate due to Heart Failure
Hospitalization Rate due to Hepatitis
Hospitalization Rate due to Hip Fractures Among Females 65+
Hospitalization Rate due to Hip Fractures Among Males 65+
Hospitalization Rate due to Hypertension
Hospitalization Rate due to Immunization-Preventable Pneumonia and Influenza
Hospitalization Rate due to Long-term Complications of Diabetes
Hospitalization Rate due to Short-term Complications of Diabetes
Hospitalization Rate due to Type 2 Diabetes
Hospitalization Rate due to Uncontrolled Diabetes
Hospitalization Rate due to Unintentional Falls
Hospitalization Rate due to Urinary Tract Infection
Emergency Room (Outpatient) Indicators
ER Rate due to Adult Asthma
ER Rate due to Asthma
ER Rate due to Asthma - Pediatric (ages 0-17)
ER Rate due to Community-Acquired Pneumonia
ER Rate due to COPD
ER Rate due to Dehydration
ER Rate due to Dental Problems
ER Rate due to Diabetes
ER Rate due to Heart Failure
ER Rate due to Hepatitis
ER Rate due to Hypertension
ER Rate due to Immunization-Preventable Pneumonia and Influenza
ER Rate due to Long-term Complications of Diabetes
ER Rate due to Short-term Complications of Diabetes
ER Rate due to Type 2 Diabetes
ER Rate due to Uncontrolled Diabetes
ER Rate due to Unintentional Falls
ER Rate due to Urinary Tract Infection

Hospitalization and ER indicators are calculated at the county and zip code levels. Data availability at the zip code level will vary depending on population density.

Behavioral Health Hospitalization and Emergency Room Indicators

Hospitalization (Inpatient) Indicators
Hospitalization Rate due to Adolescent Alcohol Use
Hospitalization Rate due to Adolescent Suicide and Intentional Self-Inflicted Injury
Hospitalization Rate due to Adult Mental Health
Hospitalization Rate due to Adult Suicide and Intentional Self-inflicted Injury
Hospitalization Rate due to Alcohol Use
Hospitalization Rate due to Opioid Use
Hospitalization Rate due to Pediatric Mental Health
Hospitalization Rate due to Substance Use

Emergency Room (Outpatient) Indicators
ER Rate due to Adolescent Alcohol Use
ER Rate due to Adolescent Suicide and Intentional Self-inflicted Injury
ER Rate due to Adult Mental Health
ER Rate due to Adult Suicide and Intentional Self-inflicted Injury
ER Rate due to Alcohol Use
ER Rate due to Opioid Use
ER Rate due to Pediatric Mental Health
ER Rate due to Substance Use

Hospitalization and ER indicators are calculated at the county and zip code levels. Data availability at the zip code level will vary depending on population density.